

Probing Confinement with Chromomagnetic Fields

Paolo Cea^a and Leonardo Cosmai^b

^aDipartimento di Fisica, Univ. of Bari and INFN - Sezione di Bari, I-70126 Bari, Italy

^bINFN - Sezione di Bari, I-70126 Bari, Italy

Using the lattice Schrödinger functional we study vacuum dynamics of SU(3) gauge theory at finite temperature. The vacuum is probed by means of an external constant Abelian chromomagnetic field. We find that by increasing the strength of the applied external field the deconfinement temperature decreases towards zero. This implies that strong enough Abelian chromomagnetic fields destroy confinement of color.

To study vacuum structure of lattice gauge theories we introduced [1,2] a gauge invariant effective action, defined by means of the lattice Schrödinger functional

$$\mathcal{Z}[U_\mu^{\text{ext}}] = \int \mathcal{D}U e^{-S_W}. \quad (1)$$

S_W is the standard Wilson action and the functional integration is extended over links on a lattice $L_s^3 \times L_4$ with the hypertorus geometry and satisfying the constraints (x_t : temporal coordinate)

$$U_k(x)|_{x_t=0} = U_k^{\text{ext}}(x), \quad (k = 1, 2, 3), \quad (2)$$

$U_k^{\text{ext}}(x)$ being the lattice version of the external continuum gauge field $\vec{A}^{\text{ext}}(x) = \vec{A}_a^{\text{ext}}(x)\lambda_a/2$.

The lattice effective action for the external static background field $\vec{A}^{\text{ext}}(x)$ is given by

$$\Gamma[\vec{A}^{\text{ext}}] = -\frac{1}{L_4} \ln \left\{ \frac{\mathcal{Z}[\vec{A}^{\text{ext}}]}{\mathcal{Z}[0]} \right\} \quad (3)$$

and is invariant for gauge transformations of the external links U_k^{ext} .

At finite temperature $T = 1/aL_t$, $L_t \ll L_s$, we introduced [3] the thermal partition function in presence of a given static background field:

$$\mathcal{Z}_T[\vec{A}^{\text{ext}}] = \int_{U_k(L_t, \vec{x})=U_k(0, \vec{x})=U_k^{\text{ext}}(\vec{x})} \mathcal{D}U e^{-S_W}. \quad (4)$$

In this case the relevant quantity is the free energy functional

$$F[\vec{A}^{\text{ext}}] = -\frac{1}{L_t} \ln \frac{\mathcal{Z}_T[\vec{A}^{\text{ext}}]}{\mathcal{Z}_T[0]}. \quad (5)$$

We used the above defined lattice effective actions to investigate the response of the vacuum to an external constant Abelian chromomagnetic field:

$$\vec{A}_a^{\text{ext}}(\vec{x}) = \vec{A}^{\text{ext}}(\vec{x})\delta_{a,3}, \quad A_k^{\text{ext}}(\vec{x}) = \delta_{k,2}x_1H. \quad (6)$$

On the lattice we have:

$$U_1^{\text{ext}}(\vec{x}) = U_3^{\text{ext}}(\vec{x}) = \mathbf{1},$$

$$U_2^{\text{ext}}(\vec{x}) = \begin{bmatrix} \exp(i\frac{gHx_1}{2}) & 0 & 0 \\ 0 & \exp(-i\frac{gHx_1}{2}) & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad (7)$$

To be consistent with the hypertorus geometry we impose that the magnetic field is quantized:

$$\frac{a^2gH}{2} = \frac{2\pi}{L_1}n_{\text{ext}}, \quad n_{\text{ext}} \text{ integer}. \quad (8)$$

This field gives rise to constant field strength, then due to gauge invariance it is easy to show that both the vacuum energy density and the free energy density are proportional to spatial volume $V = L_s^3$. Therefore the relevant quantity is the density of energy $\varepsilon[\vec{A}^{\text{ext}}]$ or free energy $f[\vec{A}^{\text{ext}}]$. At zero temperature we consider the derivative of $\varepsilon[\vec{A}^{\text{ext}}]$ with respect to β by taking gH fixed and




Figure 1. The scaling function $\kappa(\beta)$ versus β .

consider only the contribution due to the dynamical links (i.e. the links that are not constrained in the functional integration):

$$\begin{aligned} \varepsilon'_{\text{int}}[\vec{A}^{\text{ext}}] = & \left\langle \frac{1}{\Omega_{\text{int}}} \sum_{x \in \tilde{\Lambda}, \mu < \lambda} \frac{1}{3} \text{ReTr} U_{\mu\nu}(x) \right\rangle_0 \\ & - \left\langle \frac{1}{\Omega_{\text{int}}} \sum_{x \in \tilde{\Lambda}, \mu > \lambda} \frac{1}{2} \text{ReTr} U_{\mu\nu}(x) \right\rangle_{\vec{A}^{\text{ext}}} , \end{aligned} \quad (9)$$

where $\tilde{\Lambda}$ is the ensemble of the internal lattice sites. In order to extract $\varepsilon_{\text{int}}(\beta, n_{\text{ext}})$ we can numerically integrate the data for $\varepsilon'_{\text{int}}(\beta, n_{\text{ext}})$ using the trapezoidal rule.

We find that our data satisfy the scaling law:

$$x^{-\alpha} \frac{\varepsilon'_{\text{int}}(\beta, n_{\text{ext}}, L_{\text{eff}})}{\varepsilon'_{\text{ext}}} = \kappa(\beta) . \quad (10)$$

where $x = \frac{a_H}{L_{\text{eff}}}$, $a_H = \sqrt{\frac{2\pi}{gH}}$ is the magnetic length, and $L_{\text{eff}} = \Omega_{\text{int}}^{1/4}$ is the lattice effective linear size (see Fig. 1). In Eq. (10) $\varepsilon'_{\text{ext}}$ is the derivative of the classical energy due to the external applied field $\varepsilon'_{\text{ext}} = \frac{2}{3} [1 - \cos(\frac{gH}{2})] = \frac{2}{3} [1 - \cos(\frac{2\pi}{L_1} n_{\text{ext}})]$. Remarkably, the value of the exponent $\alpha = 1.5$ in Eq. (10) agrees with the one we found for the SU(2) gauge theory. From Eq. (10) we can determine the infinite volume

limit of the vacuum energy density ε_{int} :

$$\lim_{L_{\text{eff}} \rightarrow \infty} \varepsilon_{\text{int}}(\beta, n_{\text{ext}}, L_{\text{eff}}) = 0 . \quad (11)$$

As a consequence, in the continuum limit the SU(3) vacuum screens the external chromomagnetic Abelian field.

At finite temperature the relevant quantity is the β -derivative of $f[\vec{A}^{\text{ext}}]$:

$$\begin{aligned} f'[\vec{A}^{\text{ext}}] = & \left\langle \frac{1}{\Omega_{\text{int}}} \sum_{x \in \tilde{\Lambda}, \mu < \nu} \frac{1}{3} \text{ReTr} U_{\mu\nu}(x) \right\rangle_0 \\ & - \left\langle \frac{1}{\Omega_{\text{int}}} \sum_{x \in \tilde{\Lambda}, \mu < \nu} \frac{1}{3} \text{ReTr} U_{\mu\nu}(x) \right\rangle_{\vec{A}^{\text{ext}}} . \end{aligned} \quad (12)$$

It turns out that, if we start with the SU(3) gauge system at zero temperature in a constant Abelian chromomagnetic background field of fixed strength and increase the temperature, then the perturbative tail of the β -derivative of the free energy density increases with $1/L_t$ and tends towards the “classical” value. So that, we may conclude that by increasing the temperature there is no screening effect in the free energy density confirming that the zero-temperature screening of the external field is related to the confinement. Moreover, $f'_{\text{int}}(\beta, n_{\text{ext}})$ can be used to get an estimate of the deconfinement temperature T_c . To this end, we evaluate $f'[\vec{A}^{\text{ext}}]$ as a function of β for different lattice temporal sizes L_t . To determine the pseudocritical couplings we parametrize $f'(\beta, L_t)$ near the peak as

$$\frac{f'(\beta, L_t)}{\varepsilon'_{\text{ext}}} = \frac{a_1(L_t)}{a_2(L_t)[\beta - \beta^*(L_t)]^2 + 1} . \quad (13)$$

Once determined $\beta^*(L_t)$ we estimate the deconfinement temperature as

$$\frac{T_c}{\Lambda_{\text{latt}}} = \frac{1}{L_t} \frac{1}{f_{\text{SU}(3)}(\beta^*(L_t))} , \quad (14)$$

with

$$f_{\text{SU}(N)}(\beta) = \left(\frac{\beta}{2Nb_0} \right)^{b_1/2b_0^2} \exp \left(-\beta \frac{1}{4Nb_0} \right) ,$$

(15)

where N is the color number, $b_0 = (11N)/(48\pi^2)$, $b_1 = (34N^2)/(3(16\pi^2)^2)$.

Following [4] we perform a linear extrapolation to the continuum of our data for $T_c/\Lambda_{\text{latt}}$ (see Fig. 2). We see that the continuum limit deconfinement critical temperature does depend on the applied Abelian chromomagnetic field. Therefore




Figure 2. $T_c/\Lambda_{\text{latt}}$ versus aT_c for two different values of the external field strength.

we decided to vary the strength of the applied external Abelian chromomagnetic background field to study quantitatively the dependence of T_c on gH . In Fig. 3 we display our determination of T_c as a function of the external chromomagnetic field gH . We see that the critical temperature decreases by increasing the external Abelian chromomagnetic field. If the magnetic length $a_H \sim 1/\sqrt{gH}$ is the only relevant scale of the problem, then for dimensional reasons one expects that $T_c^2 \sim gH$. Indeed we try to fit our data with

$$\frac{T_c(gH)}{\Lambda_{\text{latt}}} = \frac{T_c(0)}{\Lambda_{\text{latt}}} + \alpha\sqrt{gH}. \quad (16)$$

We find a satisfying fit with $\alpha = -42.4 \pm 7.4$ and $T_c(0)/\Lambda_{\text{latt}}$ that agrees within errors with Ref. [4].




Figure 3. The continuum critical temperature $T_c/\Lambda_{\text{latt}}$ versus the external field strength gH . Solid line is the fit of Eq. (16) to our data.

We see that there exists a critical field $gH_c \simeq 0.68$ such that $T_c = 0$ for $gH > gH_c$.

The existence of a critical chromomagnetic field is naturally explained if the vacuum behaves as an ordinary relativistic color superconductor, a condensate of a color charged scalar field whose mass is proportional to the inverse of the magnetic length. A natural candidate for such tachyonic color charged scalar field is the famous Nielsen-Olesen unstable mode. However, it must be observed that the chromomagnetic condensate cannot be uniform due to gauge invariance of the vacuum, which disorders the system in such a way that there are not long range color correlations. Another important aspect of this work might be related to astrophysics applications. Our results implies the exciting possibility of a new class of compact quark stars.

REFERENCES

1. P. Cea, L. Cosmai and A.D. Polosa, Phys. Lett. B392 (1997) 177, hep-lat/9601010.
2. P. Cea and L. Cosmai, Phys. Rev. D60 (1999) 094506, hep-lat/9903005.
3. P. Cea and L. Cosmai, JHEP 11 (2001) 064.
4. J. Fingberg, U. Heller and F. Karsch, Nucl. Phys. B392 (1993) 493, hep-lat/9208012.